



## King's Research Portal

DOI:

[10.1016/j.socscimed.2016.07.004](https://doi.org/10.1016/j.socscimed.2016.07.004)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Meier, H. C. S., Haan, M. N., Mendes de Leon, C. F., Simanek, A. M., Dowd, J. B., & Aiello, A. E. (2016). Early life socioeconomic position and immune response to persistent infections among elderly Latinos. *Social Science & Medicine*, 166, 77-85. <https://doi.org/10.1016/j.socscimed.2016.07.004>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Title:** Early life socioeconomic position and immune response to persistent infections among elderly Latinos

**Authors:**

Helen C.S. Meier, PhD, MPH<sup>a</sup>, Mary N. Haan, PhD<sup>b</sup>, Carlos F. Mendes de Leon<sup>c</sup>, PhD, Amanda M. Simanek, PhD, MPH<sup>d</sup>, Jennifer B. Dowd, PhD<sup>e</sup>, Allison E. Aiello, PhD, MS<sup>f\*</sup>

**Author Affiliation:**

a. Epidemiology Branch, National Institute of Environmental Health Sciences, 111 T.W. Alexander Dr., P.O. Box 12233, MD A3-05, Research Triangle Park, NC 27709; helen.meier@nih.gov

b. Department of Epidemiology and Biostatistics, University of California San Francisco, 550 15<sup>th</sup> Street, San Francisco, CA 94158; mary.haan@ucsf.edu

c. Center for Social Epidemiology and Population Health, Department of Epidemiology, University of Michigan School of Public Health, 1415 Washington Heights, Ann Arbor, MI 48109; cmendes@umich.edu

d. Joseph J. Zilber School of Public Health, University of Wisconsin-Milwaukee, P.O. Box 413, Milwaukee, WI 53201; simaneka@uwm.edu

e. Department of Epidemiology and Biostatistics, CUNY School of Public Health, Hunter College, City University of New York, 2180 Third Ave., New York, NY 10035; jdowd@hunter.cuny.edu

f. Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, 135 Dauer Dr. 2101B McGavran-Greenberg Hall, CB 7435, Chapel Hill, NC 27599; aaiello@email.unc.edu; T: 919-966-2149, F: 919-966-2089, \*Corresponding Author

1   **Acknowledgments:**

2   Funding for the SALSA study was aided by a grant from the National Institute on Aging,  
3   National Institutes of Health: R01AG012975. Additional support for this work from grants:  
4   P60MD002249, R01DA022702, and R01DK087864. We thank Brisa Sanchez for her statistical  
5   consultation and the Stanley Neurovirology Laboratory at Johns Hopkins University, particularly  
6   Drs. Robert Yolken and Fuller Torrey, for performing the laboratory assays. This research was  
7   supported in part by the Intramural Research Program of the NIH, National Institute of  
8   Environmental Health Sciences.

9  
10   **Ethics Statement**

11   The Sacramento Area Latino Study on Aging (SALSA) was approved by the Institutional  
12   Review Board at the University of Michigan and the University of California at San Francisco  
13   and Davis.

## Abstract

Persistent infections, such as cytomegalovirus (CMV), herpes simplex virus-1 (HSV-1), *Helicobacter pylori* (*H. pylori*), and *Toxoplasma gondii* (*T. gondii*), are common in the U.S. but their prevalence varies by socioeconomic status. It is unclear if early or later life socioeconomic position (SEP) is a more salient driver of disparities in immune control of these infections. Using data from the Sacramento Area Latino Study on Aging, we examined whether early or later life SEP was the strongest predictor of immune control later in life by contrasting two life course models, the critical period model and the chain of risk model. Early life SEP was measured as a latent variable, derived from parental education and occupation, and food availability. Indicators for SEP in later life included education level and occupation. Individuals were categorized by immune response to each pathogen (seronegative, low, medium and high) with increasing immune response representing poorer immune control. Cumulative immune response was estimated using a latent profile analysis with higher total immune response representing poorer immune control. Structural equation models were used to examine direct, indirect and total effects of early life SEP on each infection and cumulative immune response, controlling for age and gender. The direct effect of early life SEP on immune response was not statistically significant for the infections or cumulative immune response. Higher early life SEP was associated with lower immune response for *T. gondii*, *H. pylori* and cumulative immune response through pathways mediated by later life SEP. For CMV, higher early life SEP was both directly associated and partially mediated by later life SEP. No association was found between SEP and HSV-1. Findings from this study support a chain of risk model, whereby early life SEP acts through later life SEP to affect immune response to persistent infections in older age.

1    **Keywords:** socioeconomic position; persistent infections; life course epidemiology; Latino  
2    health  
3

## 1   **Introduction**

2           Lower socioeconomic position (SEP) is linked to poorer individual health outcomes,  
3   though the mechanisms underlying this association are unclear (Adler & Rehkopf, 2008).  
4   Seropositivity and immune response to persistent infections may be one important pathway by  
5   which SEP influences adult health (Roberts et al., 2010; Simanek et al., 2009). Individuals of low  
6   SEP are more likely to acquire persistent infections earlier in life and are more likely to be  
7   seropositive and to have poor immune control (i.e. higher circulating antibody levels) of these  
8   pathogens across the life course (Colugnati et al., 2007; Dowd & Aiello, 2009; Dowd et al.,  
9   2009). Common persistent infections, such as cytomegalovirus (CMV), herpes simplex-1 (HSV-  
10   1), *Helicobacter pylori* (*H. pylori*), and *Toxoplasma gondii* (*T. gondii*), have been implicated in  
11   multiple chronic diseases, including cardiovascular disease, cognitive impairment, frailty, and  
12   mortality (Aiello et al., 2006; Draborg et al., 2013; Esposito et al., 2014; Hasni, 2012; Itzhaki et  
13   al., 2004; Jeon et al., 2012; Kang et al., 2004; Schmalz et al., 2005; Sorlie et al., 2000; Zajacova  
14   et al., 2009). Taken together, the social patterning of these infections may play an important role  
15   in perpetuating socioeconomic disparities in chronic diseases in older age. Whether early life  
16   SEP or later life SEP is most important in terms of shaping immune control of persistent  
17   infections and cumulative immune response to multiple infections later in life has not yet been  
18   elucidated.

19           There are several life course models by which early life SEP may be operating to  
20   influence immune response to persistent infections during adulthood. Two models, the critical  
21   period model and the chain of risk model, are strong contenders for explaining how early life  
22   SEP influences later life immune response to infection. The critical period model suggests that  
23   an event early in life, acting during a specific period of time, has a lasting independent effect on

1 later life health (Kuh, 2004). Under this model, early life SEP may serve as a critical period  
2 exposure altering an individual's susceptibility to persistent infections as well as their likelihood  
3 of exposure due to unfavorable physical and social environments, directly impacting which  
4 infections one acquires early in life and shaping their likelihood of seropositivity and immune  
5 response to the infection as an adult (Cohen et al., 2004). Increased susceptibility and frequent  
6 exposure may not only lead to infection at a younger age among individuals of low SEP in early  
7 life, but also may influence the development of the immune system by leaving a lasting early life  
8 imprint on adult immunological function (McDade, 2005). For persistent infections, which once  
9 acquired are never cleared from the body, impaired immune control later in life may manifest as  
10 increased production of antibodies targeted against these infections and are reflective of impaired  
11 cell-mediated function (Glaser & Kiecolt-Glaser, 1994).

12         Alternatively, the chain of risk model posits that exposures are linked over the life course  
13 (i.e. one experience may lead to another) to influence later life health (Kuh, 2004). Under this  
14 model, early life SEP may indirectly affect immune response to persistent infections later in life  
15 by acting through later life socioeconomic exposures. Early life social disadvantage is linked to  
16 subsequent SEP in adulthood, such as educational attainment, employment grade and income  
17 level (Oakes & Kaufman, 2006). Resources and environments corresponding to SEP over the life  
18 course may, similar to early life SEP, influence exposure to persistent infections as well as  
19 contribute to immune function and the ability to maintain immunologic control over persistent  
20 infections. For this reason, low SEP in early life may influence later life immune function  
21 indirectly by setting up a chain of risk leading to low SEP in later life which then contributes to  
22 poor immune function later in life. Thus, in a chain of risk model, later life SEP may mediate the  
23 effect of early life SEP on immune response later in life.

Limiting our ability to more broadly understand how the life course influences infectious disease outcomes, most existing studies of the social patterning of persistent infections are restricted to cross-sectional analyses in which only early life or adulthood SEP was assessed (Dowd et al., 2009; Dowd et al., 2012; Staras et al., 2008; Zajacova et al., 2009). Of those studies that have looked at the early life environment, one study suggested that adversity during childhood may negatively influence the immune system, resulting in poorer control of CMV in later life (Fagundes et al., 2013). Another study showed an association between poorer family interpersonal environment during childhood and increased later life immune response to CMV (Janicki-Deverts et al., 2014). While these studies lend support for a role of early life environment in later life immunological control, they are limited to few infections, do not directly measure early life SEP, and no study has examined potential life course models by which early life SEP influences later life immunological response.

Building upon previous work, this study addresses important gaps in our understanding of the life course SEP influences on later life immune response to multiple persistent infections. Applying a life course framework, we contrast the critical period and chain of risk models to identify the most likely pathway by which early life SEP may be operating to influence immune response to individual persistent infections and cumulative immune response to multiple infections during adulthood using data on older U.S. Latinos from the Sacramento Area Latino Study on Aging (SALSA).

## **Methods**

### *Sample*

The Sacramento Area Latino Study on Aging (SALSA) is a longitudinal cohort study of 1,789 of Mexican Americans living in the Sacramento, California metropolitan area who were



60-101 years old at baseline in 1998-1999. Detailed information about the recruitment and study population has been described previously (Haan et al., 2003). The overall response rate was 85% and more than 89% of eligible households enumerated had a least one person who participated in the study, making SALSA highly representative of older Hispanics living in Sacramento and the surrounding suburban and rural counties (Haan et al., 2003).

Data for this study come from the baseline in-home interview. To be included in the analytic sample, SALSA participants could not have missing data for covariates (age and sex, N=1,779). Using full information maximum likelihood methods, participants who had data for at least one of the six early life SEP indicators or available infection data resulting were included, resulting in a final sample of 1,562 (87.3%) for the individual infection analyses. The final sample for the cumulative immune response analysis was limited to individuals with at least one early life SEP indicator AND infection data for all pathogens (N=1,263, 70.8%).

### *Exposures*

SALSA participants provided retrospective SEP information for three time points in their life course, early life, midlife and late life. First, early life SEP was constructed as a latent variable by using five different indicators used in previous SALSA studies of early life SEP recalled by the participant and meant to capture the resources and social and physical environment: father's education (reference, fixed to 1.0), mother's education, father's occupation, mother's occupation, and food availability as a child. Parental education was measured in years. Parental occupation was measured by a 3-level categorical variable (technical, professional or managerial [high]; sales, administrative support or military [middle]; and services, manual or housewives [low]) and treated as an ordered hierarchical variable. Food availability during childhood was ascertained using the question, "When growing up, how often

1 did you not have enough to eat?” with Likert item responses and also treated as a continuous  
2 variable. Next, midlife SEP was measured by years of education completed by the SALSA  
3 participant, which is a proxy for economic resources and human capital of the participant in  
4 adulthood. Last, to measure late life SEP, the reported occupation worked by the SALSA  
5 participant for most of their life was categorized as a 4-level ordinal variable (technical,  
6 professional or managerial; sales, administrative support or military; services or manual; or  
7 housewife). Lifetime occupation is representative of later life SEP as it reflects differences in  
8 social status, earning ability and prestige.

## 9 *Outcomes*

### 10 Immune Response

11 Immune response was measured by IgG antibody level values for four pathogens, CMV,  
12 HSV-1, *H. pylori*, and *T. gondii*. Details about IgG as a serum biomarker are available in  
13 Appendix A. CMV is a beta herpesvirus and ubiquitous worldwide (Bate et al., 2010). CMV has  
14 co-evolved with humans over time and as a result has developed many mechanisms for immune  
15 evasion (Miller-Kittrell & Sparer, 2009). HSV-1 is a highly common alpha herpesvirus with  
16 adult prevalence ranging from 50 to more than 90% (Strauss & Strauss, 2008). Infections with  
17 HSV-1 are common and are characterized by recurrent oral lesions on the lips, mouth and gums  
18 (Schillinger et al., 2004). *T. gondii* is an obligate parasite that infects intestinal epithelial cells  
19 and establishes latency with intracellular bradyzoite cysts in muscle and brain cells (Galvan-  
20 Ramirez et al., 2010; Jones et al., 2014). *T. gondii* is also found worldwide with prevalence in  
21 adult populations ranging from 10% - 80% (Yolken & Torrey, 2008). *H. pylori* is a bacterium  
22 that colonizes the mucosal layer of the gastric epithelium (stomach) and generates a state of  
23 chronic inflammation (Lina et al., 2014). Persistent infection with *H. pylori* causes gastritis,

1 peptic ulcer disease and is associated with other gastric conditions, such as cancer (Lina et al.,  
2 2014).

3 Serum and plasma samples from baseline were tested at the Stanley Neurovirology  
4 Laboratory at Johns Hopkins University School of Medicine using high-throughput solid-phase  
5 enzyme-linked immunosorbent assays (ELISA) to detect pathogen-specific IgG antibody levels.  
6 The ELISA methods have been described previously (Yolken et al., 2011). Briefly, diluted  
7 aliquots of serum were reacted with antigen bound to a solid-phase surface. Quantitation of IgG  
8 for each pathogen was determined by reaction of bound antibodies with enzyme labeled anti-  
9 human IgG and enzyme substrate and optical densities were read by spectrophotometric  
10 instrumentation (Dickerson et al., 2003). The sensitivity (Sn) and specificity (Sp) of the assays  
11 are as follows: CMV Sn: 97% Sp: 94%; HSV-1 Sn: 100% Sp: 96%; *T. gondii* Sn: 94%, Sp: 96%;  
12 *H. pylori* Sn: 96% Sp: 96%.

13 Continuous antibody level was categorized by first identifying those who were  
14 seropositive and seronegative to each infection. Seropositivity was determined by standard  
15 cutoffs for each assay such that optical density unit (ODU) values less than 1.1 were classified as  
16 seronegative and values 1.1 or greater were seropositive. Among those seropositive for a given  
17 infection, continuous antibody level values were divided into tertiles and categorized as low,  
18 middle and high antibody level with those in the highest tertile representing those with the worst  
19 immune control of the infection. Based upon both pathogen serostatus and IgG antibody  
20 response we constructed a four-level hierarchical variable representing overall immune response  
21 to each pathogen in which individuals were categorized as: seronegative, low, middle and high  
22 antibody response.

## Cumulative immune response

Cumulative immune response to multiple infections may have a greater influence on disease processes than immune response towards any one infection. Recent studies suggest that traditional summary measures of cumulative immune response such as the total number of pathogens for which individuals are seropositive or exhibit elevated immune response, may not adequately capture the relative importance of particular pathogens on health outcomes of interest (Simanek et al., 2014). For this reason, we constructed a measure of cumulative immune response from a nominal latent class variable determined by the grouping of CMV, HSV-1, *H. pylori*, and *T. gondii* immune response levels. Specifically, a latent profile analysis (LPA) measurement model from continuous IgG level (log transformed) with the four infections as indicators generated latent class membership, as detailed in Appendix B. A three class solution was selected with membership as follows: the low cumulative immune response group (N=55, 4%) was seronegative to CMV and *T. gondii* and had low antibody levels to HSV-1 and *H. pylori*, the middle cumulative immune response group (N=777, 61%) was seronegative to *T. gondii* and had moderate CMV, HSV-1 and *H. pylori* antibody levels, and the high cumulative immune response group (N=431, 34%) had moderate antibody levels to all infections. Class membership was not further adjusted for potential measurement misclassification due to high entropy (Asparouhov & Muthen, 2014; Feingold et al., 2014). Predicted class membership from this LPA model was then used as a nominal dependent variable in subsequent analyses.

## *Covariates*

Covariates of interest included age and gender. Baseline age of participants was measured in years. Participant gender was categorized as male or female, and male was treated

as the referent category. Nativity was also adjusted for in sensitivity analyses with U.S. born as the referent category.

### *Statistical Analyses*

Means, percentages and standard deviations for all variables were calculated using SAS version 9.3 (SAS Institute, Inc., Cary, NC). Figure 1 depicts the theoretical life course model connecting early life SEP to late life immune response. The early life SEP latent variable was used to test three different pathways, 1) direct, independent association of early life SEP on immune response (pathway A), 2) indirect effect of early life SEP on immune response mediated by midlife SEP alone (pathway B-C), and 3) indirect effect of early life SEP on immune response mediated by midlife and later life SEP (pathway B-D-E). We did not include a pathway from early life SEP directly to later life SEP (life-long occupation) as any effect was likely mediated by midlife SEP (education). Age and gender were then added to the model with paths extending to each SEP measurement as well as the outcome to control for potential confounding.

Structural equation modeling (SEM) was used to examine the life course pathways linking early life SEP to adult antibody level for CMV, HSV-1, *H. pylori*, *T. gondii* and cumulative immune response. SEM is comprised of a measurement model and a structural model. A measurement model for the latent variable early life SEP was built using five indicators, father's education and occupation, mother's education and occupation, and food availability. The early life SEP latent variable was then used in two separate structural pathway models to predict 1) ordinal immune response (i.e. seronegative, low, middle or high antibody response) for each of the four infections and 2) cumulative immune response class while adjusting for covariates.

All SEM analyses were conducted in MPlus version 7.2 (Methuén & Methuén, Los Angeles, CA). A probit link function, theta parameterization and weighted least squares estimator (WLSMV) were used to appropriately model ordered categorical outcomes and categorical mediators for the four separate infections (Muthen, 2012). Multinomial logistic regression with Monte Carlo integration was used for the cumulative immune response model with the maximum likelihood (MLR) estimator to produce parameter estimates and standard errors that were robust to non-normality (Muthen, 2012). Goodness-of-fit for the final models was obtained using the root mean squared error of approximation (RMSEA) and comparative fit index (CFI). Models with a CFI above 0.90 and an RMSEA of less than or equal to 0.05 were considered to fit well. Model goodness-of-fit measures were not available for the cumulative immune response analysis due to the use of Monte Carlo integration.

Based on previous imputation procedures we consider the data in these analyses to be missing at random (MAR) and that age, gender and education likely explained the missing data (Haan et al., 2011; Zeki Al Hazzouri et al., 2011). Age and gender may have influenced the ability or desire to give a biospecimen during data collection and consequently the individuals for whom we had infection data. Age may also be related to recall of early life SEP. Education level commonly predicts missing information (Acock, 2013). Since age, gender and education were all included in the final models, we did not include any additional auxiliary variables to meet MAR criteria (Acock, 2013). Sensitivity analyses were performed using traditional regression and mediation techniques and similar effects sizes to the SEM results were observed (Hayes, 2013).

As a sensitivity analysis, we assessed nativity as a potential confounder, assuming nativity was associated with immune response (through some unspecified mechanism independent of SEP) and was also associated with SEP. Potential mechanisms by which nativity may influence immune response include nutrition and dietary behavior differences by place of birth. We examined whether nativity was associated with early life SEP and immune function, respectively, for each infection. Potential confounding by place of birth was only indicated for *T. gondii*, therefore, the SEM was re-run additionally controlling for nativity.

## Results

Baseline descriptive statistics of individuals included in this analysis are shown in Table 1. The sample had a mean age of 70 years and 58% were female. A total of 84% were seropositive for CMV, 91% for *H. pylori*, 87% for HSV-1, and 35% for *T. gondii*.

### *Measurement Models*

Unstandardized and standardized factor loadings for the early life SEP measurement model are presented in Supplemental Table 1. For early life SEP, all five indicators were statistically significant at  $p < 0.05$  and the measurement model fit was appropriate (RMSEA=0.05, CFI=0.94). The LPA measurement model for cumulative immune response from the four infection indicators produced a three class solution detailed in Appendix B, Supplemental Table 2 and Supplemental Table 3.

### *SEM Models: Direct, Indirect and Total Effects of Early Life SES*

A general SEM path diagram is depicted in Figure 2 for immune response to each infection and in Figure 3 for cumulative immune response. Direct, indirect and total effects of early life SEP on each of the four infection outcomes, CMV, HSV-1, *T. gondii*, and *H. pylori* are summarized in Table 2. All models fit the data well. The pathway connecting early life SEP to

immune response adjusting for all other pathways and covariates was not statistically significant in final models for any of the four infections. Early life SEP was found to indirectly influence immune response through pathways mediated by later life SEP for *T. gondii* and *H. pylori*. No association was found between early life SEP and immune response to HSV-1.

Early life SEP was associated with immune response to CMV when all pathways, direct and indirect, were considered collectively (model fit indices: RMSEA=0.031, CFI=0.956). For this total effect, a one standard deviation increase in early life SEP (one standard deviation is approximately 4 years of father's education) decreased the mean of the underlying variable for CMV antibody level by 0.11 standard deviations holding all covariates constant ( $p=0.02$ ).

The effect of early life SEP on *T. gondii* immune response was mediated by later life SEP pathways such that a one standard deviation increase in early life SEP (4 years of father's education) indirectly decreased the mean of the underlying variable for *T. gondii* antibody level by 0.10 standard deviations holding all covariates constant ( $p=0.003$ , model fit indices: RMSEA=0.031, CFI=0.962). When this indirect effect was decomposed into the education pathway and the education-occupation pathway, education was found to drive this association ( $p=0.002$ ). Further, the pathway from education directly to *T. gondii* immune response was statistically significant; a one standard deviation increase in education, 5.35 years, decreased the mean of the underlying variable for *T. gondii* antibody level by 0.21 standard deviations, holding all else constant ( $p=0.001$ ). Sensitivity analysis indicated that nativity may be a potential confounder of the SEP – *T. gondii* immune response association. When we additionally controlled for nativity in the SEM, effect estimates were attenuated and the indirect effect of early life SEP through the education pathway and the direct effect of education became marginally significant ( $p=0.07$  and  $p=0.06$  respectively) as shown in Supplemental Table 4.



Early life SEP was associated with immune response to *H. pylori* via the education-occupation pathway. A one standard deviation increase in early life SEP decreased the mean of the underlying variable for *H. pylori* antibody level by 0.03 standard deviations, holding all else constant ( $p=0.005$ , model fit indices: RMSEA=0.032, CFI=0.946). The pathway from lifetime occupation to *H. pylori* antibody level was statistically significant such that a one category increase in occupation decreased the mean of the underlying variable for *H. pylori* level by 0.11 standard deviations, holding all else constant ( $p=0.004$ ).

Table 3 shows the direct, indirect and total effects of the early life SEP latent variable on cumulative immune response. In the multinomial logistic regression model, comparing the lowest cumulative immune response group to the highest cumulative immune response group and the middle cumulative immune response group to highest cumulative immune response group, no statistically significant independent direct effect of early life SEP was found on cumulative immune response class. When considering pathways mediated by later life SEP, small, statistically significant associations were observed. Through all indirect pathways, for a one unit increase in early life SEP, the odds of being in the lowest cumulative immune response group (low immune response to HSV-1 and *H. pylori* only) were 1.11 times the odds of being in the highest cumulative immune group (moderate immune response to all infections), holding covariates constant ( $p=0.04$ ). This total indirect effect was decomposed into an indirect pathway via education only and an indirect pathway via education and lifetime occupation. The pathway via education only was shown to drive this association (OR=1.10,  $p=0.04$ ). In addition, education was found to directly influence cumulative immune response class for both the lowest v. highest cumulative immune response (OR=1.09,  $p=0.05$ ) and middle v. highest cumulative immune response (OR=1.06,  $p=0.01$ ) comparisons.

## Discussion

We examined the association between life course SEP and immune response to persistent infections and cumulative immune response. Two life course models, the critical period and the chain of risk, were evaluated to determine how life course SEP impacts immune response to four persistent infections associated with chronic health conditions later in life. Early life SEP was not independently associated with immune response in older age for any of the four persistent infections included in this study or the cumulative immune response to these infections. Instead, higher later life SEP, as measured by education and lifetime occupation, mediated the effect of early life SEP on immune response. These findings support a chain of risk model, whereby early life SEP influences later life SEP, which in turn, impacts immune control of these infections.

Education, in particular, may be a key link in the socioeconomic chain of risk. This may be due to the far-reaching influence of higher educational attainment on the potential for subsequent higher SEP exposures, such as higher occupational status, higher income level and increased wealth. Higher education also contributes to the development of health capital and better health behaviors, potentially reducing individual susceptibility or improving immunological function through, for example, better nutrition. In addition, higher educational attainment and higher later life SEP may act as a buffer by reducing the number and severity of stressors, which may impact the frequency of reactivation to persistent infections. Therefore, early life conditions may set the stage to enhance or prevent educational achievement, which has life-long consequences for social advantage or disadvantage, influences disparities in immune response to persistent infections, and general health overall. These findings have key public health implications because higher adult immune response to these infections has been associated with mortality and chronic health conditions (Aiello et al., 2009; Aiello et al., 2006; Dickerson et

1 al., 2003; Esposito et al., 2014; Simanek et al., 2009). Moreover, there are significant disparities  
2 in these infections by race/ethnicity, indicating that minority populations are more likely to  
3 experience detrimental impacts of poor life course SEP on immune response which may affect  
4 overall health and increased mortality.

5 Of the four persistent infections examined in this study, early life SEP may be most  
6 important for later life immune response to CMV. Early life SEP was inversely associated with  
7 CMV immune response after adjusting for covariates but not mediators. Though the independent  
8 effect of early life SEP controlling for all mediating pathways was not statistically significant,  
9 the role early life may play for acquiring CMV and subsequent immune control of this infection  
10 is not diminished. CMV is generally acquired early in life through direct contact with infected  
11 body fluids (Centers for Disease Control and Prevention, 2010). Cross-sectional evidence shows  
12 racial and socioeconomic disparities exist for CMV seroprevalence and IgG antibody levels that  
13 begin at young ages (Dowd et al., 2012). In addition, immune response to CMV increases with  
14 age and lower SEP (Dowd & Aiello, 2009), and Mexican Americans on average acquire CMV  
15 10 years earlier than non-Hispanic Whites (Colugnati et al., 2007). Low SEP in early life may  
16 result in earlier infection with CMV and fewer resources available to manage the infection,  
17 increasing the frequency of reactivation over the life course. More frequent reactivations may  
18 produce negative downstream biological effects, including inducing inflammation and the  
19 exhaustion of cell mediated immunity (immunosenescence) (Pawelec et al., 2006).

20 Our study findings also suggest there may be evidence for a “trigger effect” for immune  
21 response to *T. gondii* and *H. pylori*. A trigger effect occurs in a chain of risk where only the final  
22 link in the chain has a marked effect on the outcome (Ben-Shlomo & Kuh, 2002). We observed  
23 a direct effect of adulthood education on immune response to *T. gondii* independent of all other

1 pathways. This may mean that educational attainment is a trigger for immune control of *T.*  
2 *gondii* later in life. The direct effect of education and the indirect effect of early life SEP through  
3 the education pathway was attenuated and made marginally significant by additionally  
4 controlling for nativity. Of note, *T. gondii* is transmitted through cat feces, under cooked meat,  
5 contaminated water or may be inhaled or ingested directly through soil (Torrey & Yolken, 2013).  
6 It is therefore possible that *T. gondii* immune response may differ by country, if these potential  
7 exposure routes are unrelated to socioeconomic differences. Better education may not only  
8 prevent exposure to *T. gondii* through knowledge of these transmission pathways, but it also may  
9 represent better access to other resources, such as health care, which influences overall health  
10 and immune system function later in life.

11 Similarly, we observed a direct effect of lifetime occupation on immune response to *H.*  
12 *pylori* independent of all other pathways and that early life SEP was only associated with this  
13 outcome through the education-occupation pathway. Occupation may, therefore, be a trigger for  
14 immune control of *H. pylori* later in life. This is consistent with previous studies, which have  
15 shown that manual or unskilled workers have higher odds of *H. pylori* infection than non-manual  
16 workers (Moayyedi et al., 2002; Rosenstock et al., 1996). Lower occupational levels may  
17 increase the virulence of *H. pylori* as well as interfere with the immune response to infection  
18 through increased energy expenditure or other pathways (Rosenstock et al., 1996). In addition, a  
19 recent study found that the inability for employees to make decisions affecting the experience of  
20 fairness was associated with increased long-term levels of inflammatory markers CRP and IL-6  
21 among men (Elovainio et al., 2010). A systematic review of psychosocial job stress and  
22 immunity found that greater job satisfaction may have a positive impact on immune outcomes  
23 and that unemployment and job security are significant factors leading to reduced cellular

immune response (Nakata, 2012). Thus, lower occupational status may influence immunological control of *H. pylori*, potentially through inflammatory pathways, and result in decreased resources to handle stress and its negative health consequences. Alternatively, jobs that do not provide health care coverage may result in fewer doctor visits associated with *H. pylori* infection. *H. pylori* is associated with peptic ulcer disease and can be treated with a combination of proton pump inhibitors and antibiotics (Grad et al., 2012). Our study did not directly diagnose *H. pylori* infection but IgG antibody levels to the infection indicate an initial or lasting immune response to the infection once it has cleared (Grad et al., 2012). Treatment for *H. pylori* or other incidental antibiotic use may result in seroreversion or decreased measurable antibody levels (Grad et al., 2012). As a result, higher SEP and health care access due to employment with health care benefits may result in lower immune response to *H. pylori*.

Education was also found to mediate the effect of early life SEP on cumulative immune response; those with higher education were more likely to have better cumulative immune control. Lower SEP during early life and midlife may impact individual susceptibility and patterns of exposure to these infections, as well as immunological control over the life course. Children with lower SEP may have worse housing conditions and poorer hygiene than children with higher SEP, increasing the likelihood of exposure and acquisition of multiple persistent infections. Further, co-infection may not only increase the risk of infection to other pathogens, but also increase the severity of subsequent infections (Espinola-Klein et al., 2002; Rubicz et al., 2011). Indeed, recent studies provide convincing evidence that the measles virus reduces immune response after initial infection leading to heightened susceptibility to other infections and this may explain why childhood measles vaccination led to dramatic reductions in infections with other pathogens in childhood (Griffin, 2010). Likewise, seropositivity to multiple persistent

1 infections may contribute to morbidity over and above the influence of each infection alone  
2 through additional tolls placed on the immune system and activation of inflammatory pathways  
3 (Aiello et al., 2009). These results reinforce the importance of education for reducing exposure to  
4 persistent infections and suggest improving educational attainment as a potential target for public  
5 health intervention aimed at reducing disparities in these infections. Other potential targets may  
6 include the development of vaccinations or treatments for the pathogens that do not currently  
7 have either of these public health or medical measures.

8         This study has some limitations. The SALSA study may be subject to survivor bias  
9 because participants were at least 60 years of age at baseline; participants may be healthier and  
10 have better control of persistent infections than those who did not survive to enrollment, biasing  
11 our results towards the null. In addition, early life SEP measures are prone to recall bias.  
12 However, reporting of early life SEP is unlikely to vary by immune response, as individuals are  
13 unlikely to be aware of their serostatus and immunological response to infection. None of these  
14 infections are routinely diagnosed in populations unless they are associated with overt  
15 symptoms, such as peptic ulcer disease. A previous study in the SALSA cohort found that only  
16 0.8% were taking antibiotics to treat *H. pylori* infection and among those seropositive to *H.*  
17 *pylori* 11% reported taking medications for acid suppression, peptic disorders, or antacids at the  
18 time of data collection for this study (Jeon et al., 2012). In this elderly cohort, poor cognition  
19 may influence the recall of early life SEP but we expect this to be non-differential with respect to  
20 level of SEP, biasing estimates toward the null. As with many studies of infections, we have no  
21 information on primary infection timing. While it is possible infections acquired earlier in life  
22 may influence later life SEP attainment, this does not prevent us from examining the role of SEP  
23 over the life course in relation to immune control later in life. Further, without infection timing,

we were unable to investigate potentially protective health effects that these pathogens may have in early life (Amberbir et al., 2014; Rook, 2012). This purpose of this study was to determine the role of early life SEP in the social patterning of immune response to persistent pathogens. The clinical significance of a change in antibody level to the chronic infections examined here has no direct correlate with common measures of immune functioning (e.g. vaccine response) because there are currently no vaccinations for any of these infections. As a surrogate, we compared our results to vaccination with varicella zoster virus (VZV), which is another herpesvirus that affects children and adults. Vaccination of adults with VZV results in a geometric increase of 2.3 compared to control subjects (Gilbert et al., 2014). Interpreting our results in light of this immune response to VZV vaccine, we observed a CMV antibody level geometric mean decrease of 0.22 for those with both high early life SEP and high midlife SEP relative to all others (in the case of chronic infection, a lower antibody level is protective). This corresponds to a 10<sup>th</sup> of what one would observe when directly vaccinating an adult with VZV and is equal to a 12% lower CMV antibody level for those in the highest SEP, suggesting a potential clinically meaningful change, especially in light of the fact that we did not directly vaccinate individuals. However, this interpretation should be made with caution given that immune response to a CMV vaccine, if ever developed, may not show the same change in antibody level as VZV. VZV was not directly measured in this study because there is a vaccine available and it would be impossible to disentangle a vaccine response versus natural immune response. Natural immune response was observed for the pathogens included in this study—a strength of our work, since there are no available vaccines. It is not possible to compare surrogate vaccine responses for *H. pylori* or *T. gondii* immune response since there are no similar vaccines. Future vaccine challenge studies should consider collecting life course SEP data to assess the influence of these

factors on vaccination response in adulthood with clinically significant immune response outcomes. Finally, there are additional persistent infections for which we do not have data, however, the four that we studied are important to understand because they are associated with chronic health outcomes and more prevalent among Latino populations (Aiello et al., 2006; Hasni, 2012; Simanek et al., 2009; Simanek et al., 2011; Sorlie et al., 2000; Zhu et al., 2000).

Despite these limitations, this study has many strengths. We use a life course approach to determine how early life social environment influences immune response to persistent infections later in life in a well-characterized cohort of predominantly Mexican Americans, limiting the influences of significant heterogeneity present among Latino populations. SALSA is a large cohort with rich data both on life course SEP and immune response to persistent pathogens. Samples were tested at a reliable laboratory using validated methods. Using SEM, we maximized the use of available SEP information, incorporated measurement error into the modeling process and estimated pathway specific effects and standard errors.

This study adds to the growing literature showing that childhood may not be an independent critical period for SEP exposures in relation to later life health outcomes, and instead the beginning of multiple mediating pathways that ultimately influence adult health. Future research should expand on this work to examine the effects of nativity and acculturation, important components of social and racial/ethnic disparities in health, as independent predictors of immune response, as well as modifiers of the life course social patterning of immune response to persistent infections. In addition, other life course models, such as the social mobility model, may also fit these relations of interest and provide insight into whether later life SEP partially or entirely remediates or modifies low early life SEP exposure. Lastly, given the complex relationship between the gut microbiome, infection and immune response, future research should



consider additional microbial pathways that may be linked to SEP and ultimately affect immune response over the life course.

### **Conclusion**

Numerous studies have demonstrated that disparities exist in the seroprevalence and immune control of persistent infections by SEP beginning early in life, suggesting childhood may be an important period during the life course for acquiring and establishing control of persistent infections. Our study found that higher immune response to chronic infections later in life is not just a function of disadvantage during early life, but instead reflects the end result of a chain of socioeconomic disadvantage throughout the life span. Our findings suggest that interventions targeting increased educational attainment as a modifiable risk factor may be effective for improving immune control of persistent infections later in life. Moreover, understanding the life course SEP pathways that influence immune response to persistent infections may help shed light on why chronic disease disparities associated with these infections appear to persist by SEP over time and across generations in the U.S.

## References

- Acock, A.C. (2013). *Discovering Structural Equation Modeling Using Stat*. College Station, TX: Stata Press.
- Adler, N.E., & Rehkopf, D.H. (2008). U.S. disparities in health: descriptions, causes, and mechanisms. *Annu Rev Public Health*, 29, 235-252.
- Aiello, A.E., Diez-Roux, A., Noone, A.M., Ranjit, N., Cushman, M., Tsai, M.Y., et al. (2009). Socioeconomic and psychosocial gradients in cardiovascular pathogen burden and immune response: the multi-ethnic study of atherosclerosis. *Brain Behav Immun*, 23, 663-671.
- Aiello, A.E., Haan, M., Blythe, L., Moore, K., Gonzalez, J.M., & Jagust, W. (2006). The influence of latent viral infection on rate of cognitive decline over 4 years. *J Am Geriatr Soc*, 54, 1046-1054.
- Amberbir, A., Medhin, G., Abegaz, W.E., Hanlon, C., Robinson, K., Fogarty, A., et al. (2014). Exposure to *Helicobacter pylori* infection in early childhood and the risk of allergic disease and atopic sensitization: a longitudinal birth cohort study. *Clin Exp Allergy*, 44, 563-571.
- Asparouhov, T., & Muthen, B. (2014). Auxiliary Variables in Mixture Modeling: Three-Step Approaches Using Mplus. *Structural Equation Modeling-a Multidisciplinary Journal*, 21, 329-341.
- Bate, S.L., Dollard, S.C., & Cannon, M.J. (2010). Cytomegalovirus seroprevalence in the United States: the national health and nutrition examination surveys, 1988-2004. *Clin Infect Dis*, 50, 1439-1447.
- Ben-Shlomo, Y., & Kuh, D. (2002). A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *Int J Epidemiol*, 31, 285-293.
- Centers for Disease Control and Prevention. (2010). Cytomegalovirus and Congenital CMV Infection. In D.o.V.D. National Center for Immunization and Respiratory Diseases (Ed.).
- Cohen, S., Doyle, W.J., Turner, R.B., Alper, C.M., & Skoner, D.P. (2004). Childhood socioeconomic status and host resistance to infectious illness in adulthood. *Psychosom Med*, 66, 553-558.
- Colugnati, F.A., Staras, S.A., Dollard, S.C., & Cannon, M.J. (2007). Incidence of cytomegalovirus infection among the general population and pregnant women in the United States. *BMC Infect Dis*, 7, 71.
- Dickerson, F.B., Boronow, J.J., Stallings, C., Origoni, A.E., Ruslanova, I., & Yolken, R.H. (2003). Association of serum antibodies to herpes simplex virus 1 with cognitive deficits in individuals with schizophrenia. *Arch Gen Psychiatry*, 60, 466-472.
- Dowd, J.B., & Aiello, A.E. (2009). Socioeconomic differentials in immune response. *Epidemiology*, 20, 902-908.
- Dowd, J.B., Aiello, A.E., & Alley, D.E. (2009). Socioeconomic disparities in the seroprevalence of cytomegalovirus infection in the US population: NHANES III. *Epidemiol Infect*, 137, 58-65.
- Dowd, J.B., Palermo, T.M., & Aiello, A.E. (2012). Family poverty is associated with cytomegalovirus antibody titers in U.S. children. *Health Psychol*, 31, 5-10.

- Draborg, A.H., Duus, K., & Houen, G. (2013). Epstein-Barr virus in systemic autoimmune diseases. *Clin Dev Immunol*, 2013, 535738.
- Elovainio, M., Ferrie, J.E., Singh-Manoux, A., Gimeno, D., De Vogli, R., Shipley, M., et al. (2010). Organisational justice and markers of inflammation: the Whitehall II study. *Occup Environ Med*, 67, 78-83.
- Espinola-Klein, C., Rupprecht, H.J., Blankenberg, S., Bickel, C., Kopp, H., Rippin, G., et al. (2002). Impact of infectious burden on extent and long-term prognosis of atherosclerosis. *Circulation*, 105, 15-21.
- Esposito, S., Bosis, S., Semino, M., & Rigante, D. (2014). Infections and systemic lupus erythematosus. *Eur J Clin Microbiol Infect Dis*.
- Fagundes, C.P., Glaser, R., Malarkey, W.B., & Kiecolt-Glaser, J.K. (2013). Childhood adversity and herpesvirus latency in breast cancer survivors. *Health Psychol*, 32, 337-344.
- Feingold, A., Tiberio, S.S., & Capaldi, D.M. (2014). New Approaches for Examining Associations With Latent Categorical Variables: Applications to Substance Abuse and Aggression. *Psychology of Addictive Behaviors*, 28, 257-267.
- Galvan-Ramirez, M.L., Madriz Elisondo, A.L., Rico Torres, C.P., Luna-Pasten, H., Rodriguez Perez, L.R., Rincon-Sanchez, A.R., et al. (2010). Frequency of *Toxoplasma gondii* in pork meat in Ocotlan, Jalisco, Mexico. *J Food Prot*, 73, 1121-1123.
- Gilbert, P.B., Gabriel, E.E., Miao, X., Li, X., Su, S.C., Parrino, J., et al. (2014). Fold rise in antibody titers by measured by glycoprotein-based enzyme-linked immunosorbent assay is an excellent correlate of protection for a herpes zoster vaccine, demonstrated via the vaccine efficacy curve. *J Infect Dis*, 210, 1573-1581.
- Glaser, R., & Kiecolt-Glaser, J. (1994). Stress-associated immune modulation and its implications for reactivation of latent herpesviruses. In R. Glaser, & J. Jones (Eds.), *Human herpesvirus infections* pp. 245-270). New York: Dekker.
- Grad, Y.H., Lipsitch, M., & Aiello, A.E. (2012). Secular trends in *Helicobacter pylori* seroprevalence in adults in the United States: evidence for sustained race/ethnic disparities. *Am J Epidemiol*, 175, 54-59.
- Griffin, D.E. (2010). Measles virus-induced suppression of immune responses. *Immunol Rev*, 236, 176-189.
- Haan, M.N., Mungas, D.M., Gonzalez, H.M., Ortiz, T.A., Acharya, A., & Jagust, W.J. (2003). Prevalence of dementia in older latinos: the influence of type 2 diabetes mellitus, stroke and genetic factors. *J Am Geriatr Soc*, 51, 169-177.
- Haan, M.N., Zeki Al-Hazzouri, A., & Aiello, A.E. (2011). Life-span socioeconomic trajectory, nativity, and cognitive aging in Mexican Americans: the Sacramento Area Latino Study on Aging. *J Gerontol B Psychol Sci Soc Sci*, 66 Suppl 1, i102-110.
- Hasni, S.A. (2012). Role of *Helicobacter pylori* infection in autoimmune diseases. *Curr Opin Rheumatol*, 24, 429-434.
- Hayes, A.F. (2013). *Introduction to mediation, moderation, and conditional process analysis : a regression-based approach*.
- Itzhaki, R.F., Wozniak, M.A., Appelt, D.M., & Balin, B.J. (2004). Infiltration of the brain by pathogens causes Alzheimer's disease. *Neurobiol Aging*, 25, 619-627.
- Janicki-Deverts, D., Cohen, S., Doyle, W.J., Marsland, A.L., & Bosch, J. (2014). Childhood environments and cytomegalovirus serostatus and reactivation in adults. *Brain Behav Immun*.

- Jeon, C.Y., Haan, M.N., Cheng, C., Clayton, E.R., Mayeda, E.R., Miller, J.W., et al. (2012). *Helicobacter pylori* infection is associated with an increased rate of diabetes. *Diabetes Care*, 35, 520-525.
- Jones, J.L., Kruszon-Moran, D., Rivera, H.N., Price, C., & Wilkins, P.P. (2014). *Toxoplasma gondii* seroprevalence in the United States 2009-2010 and comparison with the past two decades. *Am J Trop Med Hyg*, 90, 1135-1139.
- Kang, I., Quan, T., Nolasco, H., Park, S.H., Hong, M.S., Crouch, J., et al. (2004). Defective control of latent Epstein-Barr virus infection in systemic lupus erythematosus. *J Immunol*, 172, 1287-1294.
- Kuh, D., Ben-Shlomo, Y., eds. (2004). *A Life Course Approach to Chronic Disease Epidemiology*. New York, NY: Oxford University Press.
- Lina, T.T., Alzahrani, S., Gonzalez, J., Pinchuk, I.V., Beswick, E.J., & Reyes, V.E. (2014). Immune evasion strategies used by. *World J Gastroenterol*, 20, 12753-12766.
- McDade, T.W. (2005). Life history, maintenance, and the early origins of immune function. *Am J Hum Biol*, 17, 81-94.
- Miller-Kittrell, M., & Sparer, T.E. (2009). Feeling manipulated: cytomegalovirus immune manipulation. *Virol J*, 6, 4.
- Moayyedi, P., Axon, A.T., Feltbower, R., Duffett, S., Crocombe, W., Braunholtz, D., et al. (2002). Relation of adult lifestyle and socioeconomic factors to the prevalence of *Helicobacter pylori* infection. *Int J Epidemiol*, 31, 624-631.
- Muthen, L.K.M.B.O. (2012). *Mplus Statistical Analysis with Latent Variables User's Guide*. Los Angeles, CA: Muthen & Muthen.
- Nakata, A. (2012). Psychosocial job stress and immunity: a systematic review. *Methods Mol Biol*, 934, 39-75.
- Oakes, J.M., & Kaufman, J.S. (2006). *Methods in Social Epidemiology*. San Francisco, CA: Jossey-Bass.
- Pawelec, G., Koch, S., Franceschi, C., & Wikby, A. (2006). Human immunosenescence: does it have an infectious component? *Ann N Y Acad Sci*, 1067, 56-65.
- Roberts, E.T., Haan, M.N., Dowd, J.B., & Aiello, A.E. (2010). Cytomegalovirus antibody levels, inflammation, and mortality among elderly Latinos over 9 years of follow-up. *Am J Epidemiol*, 172, 363-371.
- Rook, G.A. (2012). Hygiene hypothesis and autoimmune diseases. *Clin Rev Allergy Immunol*, 42, 5-15.
- Rosenstock, S.J., Andersen, L.P., Rosenstock, C.V., Bonnevie, O., & Jorgensen, T. (1996). Socioeconomic factors in *Helicobacter pylori* infection among Danish adults. *Am J Public Health*, 86, 1539-1544.
- Rubicz, R., Leach, C.T., Kraig, E., Dhurandhar, N.V., Grubbs, B., Blangero, J., et al. (2011). Seroprevalence of 13 common pathogens in a rapidly growing U.S. minority population: Mexican Americans from San Antonio, TX. *BMC Res Notes*, 4, 433.
- Schillinger, J.A., Xu, F., Sternberg, M.R., Armstrong, G.L., Lee, F.K., Nahmias, A.J., et al. (2004). National seroprevalence and trends in herpes simplex virus type 1 in the United States, 1976-1994. *Sex Transm Dis*, 31, 753-760.
- Schmaltz, H.N., Fried, L.P., Xue, Q.L., Walston, J., Leng, S.X., & Semba, R.D. (2005). Chronic cytomegalovirus infection and inflammation are associated with prevalent frailty in community-dwelling older women. *J Am Geriatr Soc*, 53, 747-754.

- Simanek, A.M., Dowd, J.B., & Aiello, A.E. (2009). Persistent pathogens linking socioeconomic position and cardiovascular disease in the US. *Int J Epidemiol*, 38, 775-787.
- Simanek, A.M., Dowd, J.B., Pawelec, G., Melzer, D., Dutta, A., & Aiello, A.E. (2011). Seropositivity to cytomegalovirus, inflammation, all-cause and cardiovascular disease-related mortality in the United States. *PLoS One*, 6, e16103.
- Simanek, A.M., Dowd, J.B., Zajacova, A., & Aiello, A.E. (2014). Unpacking the 'black box' of total pathogen burden: is number or type of pathogens most predictive of all-cause mortality in the United States? *Epidemiol Infect*, 1-11.
- Sorlie, P.D., Nieto, F.J., Adam, E., Folsom, A.R., Shahar, E., & Massing, M. (2000). A prospective study of cytomegalovirus, herpes simplex virus 1, and coronary heart disease: the atherosclerosis risk in communities (ARIC) study. *Arch Intern Med*, 160, 2027-2032.
- Staras, S.A., Flanders, W.D., Dollard, S.C., Pass, R.F., McGowan, J.E., Jr., & Cannon, M.J. (2008). Cytomegalovirus seroprevalence and childhood sources of infection: A population-based study among pre-adolescents in the United States. *J Clin Virol*, 43, 266-271.
- Strauss, E.G., & Strauss, J.H. (2008). *Viruses and Human Disease*. Boston, M.A.: Elsevier.
- Torrey, E.F., & Yolken, R.H. (2013). Toxoplasma oocysts as a public health problem. *Trends Parasitol*, 29, 380-384.
- Yolken, R.H., & Torrey, E.F. (2008). Are some cases of psychosis caused by microbial agents? A review of the evidence. *Mol Psychiatry*, 13, 470-479.
- Yolken, R.H., Torrey, E.F., Lieberman, J.A., Yang, S., & Dickerson, F.B. (2011). Serological evidence of exposure to Herpes Simplex Virus type 1 is associated with cognitive deficits in the CATIE schizophrenia sample. *Schizophr Res*, 128, 61-65.
- Zajacova, A., Dowd, J.B., & Aiello, A.E. (2009). Socioeconomic and race/ethnic patterns in persistent infection burden among U.S. adults. *J Gerontol A Biol Sci Med Sci*, 64, 272-279.
- Zeki Al Hazzouri, A., Haan, M.N., Kalbfleisch, J.D., Galea, S., Lisabeth, L.D., & Aiello, A.E. (2011). Life-course socioeconomic position and incidence of dementia and cognitive impairment without dementia in older Mexican Americans: results from the Sacramento area Latino study on aging. *Am J Epidemiol*, 173, 1148-1158.
- Zhu, J., Quyyumi, A.A., Norman, J.E., Csako, G., Waclawiw, M.A., Shearer, G.M., et al. (2000). Effects of total pathogen burden on coronary artery disease risk and C-reactive protein levels. *Am J Cardiol*, 85, 140-146.

Figure 1. Life course model of the association between early life SEP and adult immune response to common persistent pathogens




Figure 2. General diagram of SEM for indirect and direct effects of early life SEP on immune response




Figure 3. General diagram of SEM for indirect and direct effects of early life SEP on cumulative immune response






Table 1. Descriptive statistics for the Sacramento Area Latino Study on Aging (SALSA) at baseline

Variable	N	Mean or %	SD
Early Life SEP			
Father's Education (years)	732	3.35	4.18
Mother's Education (years)	827	3.37	3.12
Father's Occupation	1132		
High	88	7.80	
Middle	54	4.80	
Low	990	87.50	
Mother's Occupation	1227		
High	41	3.30	
Middle	15	1.20	
Low	1171	95.40	
Food Availability	1249	4.55	0.98
Sibling Mortality	1247	1.26	1.14
Midlife SEP			
Education (years)	1562	7.53	5.35
Late Life SEP			
Technical, Professional or Managerial	182	11.80	
Sales, Administrative Support or Military	172	11.15	
Services or Manual	907	58.82	
Housewives	281	18.22	
Covariates			
Age (years)	1562	70.34	6.93
Female	1562	58.00	0.49
Seropositive			
CMV	1263	84.00	0.36
<i>T. gondii</i>	1263	35.00	0.48
<i>H. pylori</i>	1263	91.00	0.28
HSV-1	1263	87.00	0.34
Cumulative Immune Response			
Low Group	55	4.00	
Medium Group	777	61.00	
High Group	431	35.00	

Table 2. Standardized direct and indirect effects of SEP on later life immune response, N=1562

			Persistent Infection					
	CMV	p-value	HSV-1	p-value	<i>T. gondii</i>	p-value	<i>H. pylori</i>	p-value
Early Life SEP								
Total Effect	<b>-0.108</b>	<b>0.018</b>	-0.067	0.164	0.014	0.813	-0.072	0.127
Direct Effect	-0.086	0.185	-0.077	0.254	0.115	0.165	-0.078	0.231
Total Indirect Effect	-0.023	0.405	0.01	0.725	<b>-0.101</b>	<b>0.003</b>	0.006	0.831
Decomposition of Indirect Effects								
via Education	-0.024	0.441	-0.001	0.962	<b>-0.119</b>	<b>0.002</b>	0.037	0.211
via Education and Occupation	0.001	0.949	0.011	0.301	0.067	0.142	<b>-0.031</b>	<b>0.005</b>
Midlife SEP								
Direct Effect of Education	-0.041	0.442	-0.003	0.962	<b>-0.209</b>	<b>0.001</b>	0.065	0.206
Late Life SEP								
Direct Effect of Occupation	0.003	0.949	0.042	0.299	0.067	0.142	<b>-0.114</b>	<b>0.004</b>
Model Fit								
RMSEA	0.031		0.031		0.031		0.029	
CFI	0.956		0.962		0.96		0.965	

All models adjusted for age and sex. **Bold** is statistically significant at p=0.05.

Table 3. Life course SEP effects on later life cumulative immune response

	Cumulative Immune Response			
	Low v. High (N=1263)		Middle v. High (N=1263)	
	OR	p-value	OR	p-value
Early life SEP				
Total Effect	1.17	0.06	0.83	0.11
Direct Effect	1.05	0.64	0.93	0.16
Total Indirect Effect	<b>1.11</b>	<b>0.04</b>	0.89	0.08
Decomposition of Indirect Effects				
via Education	<b>1.10</b>	<b>0.05</b>	0.92	0.16
via Education and Occupation	1.02	0.73	0.97	0.17
Midlife SEP				
Direct Effect of Education	<b>1.09</b>	<b>0.05</b>	<b>1.06</b>	<b>0.01</b>
Late Life SEP				
Direct Effect of Occupation	1.07	0.73	0.89	0.16

All models adjusted for Age and Sex

Standardized Estimates. **Bold** is statistically significant at p=0.05

## **Appendix A.**

### *Immune Response*

Lasting immune response is measured by the amount of IgG antibody to the specific pathogen circulating in the blood. IgG is a serum biological marker indicating that an individual received sufficient exposure to the pathogen to activate the adaptive immune response at some point over the life course. Recent infection is measured by IgM, however this antibody isotype is short-lived and difficult to capture in a population-based study. Moreover, recent infections with CMV, HSV-1, *H. pylori* and *T. gondii* in older age are generally rare in the U.S. as a majority of individuals are exposed at a young age (childhood-adolescence) and even earlier in Mexican Americans. Persistent infections establish residence in the body after initial infection and periodically reactivate inducing an increase in IgG antibody response. For these reasons, elevated measures of IgG levels among older age individuals are likely due to reactivation of a pathogen from exposure to stressor or age-related changes in immune competence.

## Appendix B.

### *LPA Measurement Model*

Supplementary Table 1 shows the results of the LPA measurement model for cumulative immune response from the four infection indicators for a one, two, three, four and five class solution. The best class number solution was determined based on comparison of standard fit measures, including the Bayesian Information Criterion (BIC), sample size adjusted BIC, the Lo, Mendell and Ruben log-likelihood ratio test, and the bootstrapped log-likelihood ratio test (BLRT), as well as judgment on the nature of the groups (mean antibody level) and their interpretability in relation to theory and previous research (Marsh et al., 2009). Fit statistics were used to first compare all possible solutions. The three and four class solutions emerged as suitable for the data as the Lo, Medell, Rubin LRT was not statistically significant for the five-class solution. Then, the mean IgG level values for each group were reviewed for the three class and four class solution. Upon considering the nature of the groups and their interpretability, as well as prior research, a three class solution was deemed most appropriate even though the Lo, Mendell, Rubin LRT and the BLRT suggested that the four class solution was a numerical improvement over the three class solution. Class membership for this solution is as follows: the low cumulative immune response group (N=55, 4%) was seronegative to CMV and *T. gondii* and had mean IgG antibody levels to HSV-1 and *H. pylori* in the lowest tertile, the middle cumulative immune response group (N=777, 61%) was seronegative to *T. gondii* and had mean CMV, HSV-1 and *H. pylori* IgG antibody levels in the middle tertile, and the high cumulative immune response group (N=431, 34%) had mean IgG antibody levels in the middle tertile to all infections. The mean IgG antibody level for each infection within each class group is reported in Supplementary Table 2. The three class solution had high entropy (0.88), indicating good class

separation. Predicted class membership was extracted and then used as a nominal dependent variable in the SEM analysis with high cumulative immune response as the referent category.

Supplemental Table1. Measurement model for early life SEP, model fit: RMSEA=0.05, CFI=0.94

Indicator Loadings	Unstandardized Estimate		SE	Standardized Estimate		SE
Father's Education	1.00 (fixed)			0.73 ***		0.054
Mother's Education	0.763 ***		0.113	0.652 ***		0.049
Father's Occupation	0.036 ***		0.028	0.437 ***		0.054
Mother's Occupation	0.153 ***		0.029	0.380 ***		0.071
Food Availability	0.036 *		0.015	0.117 *		0.049

\*Significant at  $p < 0.05$ , \*\*Significant at  $p < 0.01$ , \*\*\*Significant at  $p < 0.001$

Supplemental Table 2. Measurement models for cumulative immune response from LPA with log continuous IgG level indicators for four infections (n=1263)

Model Fit Measure	1 Class	2 Class	3 Class	4 Class	5 Class
AIC	10112	9747	9653	9507	9435
BIC	10153	9814	9745	9626	9579
Sample Size Adjusted BIC	10128	9773	9688	9552	9489
Entropy	n/a	0.875	0.879	0.836	0.856
		365		151	
Lo, Mendell, Rubin LRT	n/a	(p<0.001)	102 (0.02)	(p<0.001)	80 (0.06)
Bootstrapped Likelihood Ratio					
Test	n/a	p<0.001	p<0.001	p<0.001	p<0.001




Supplemental Table 3. Mean IgG levels for the cumulative immune response three class measurement model solution (n=1263)

Cumulative Immune Response	Mean IgG Level (ODU)			
	CMV	HSV-1	<i>H. pylori</i>	<i>T. gondii</i>
Low Group	0.05	1.67	2.57	0.40
Medium Group	2.02	2.10	3.15	0.47
High Group	1.85	2.16	3.24	2.37


Supplemental Table 4. Standardized direct and indirect effects of SEP on later life immune response to *T. gondii* controlling for age, sex and nativity, N=1562

	<i>T. gondii</i>	p-value
Early Life SEP		
Total Effect	0.031	0.599
Direct Effect	0.084	0.319
Total Indirect Effect	-0.053	0.14
Decomposition of Indirect Effects		
via Education	-0.07	0.071
via Education and Occupation	0.017	0.137
Midlife SEP		
Direct Effect of Education	-0.132	0.067
Late Life SEP		
Direct Effect of Occupation	0.067	0.133
Model Fit		
RMSEA	0.037	
CFI	0.953	


Supplemental Figure 1. Diagram of SEM for standardized direct and indirect effects of early life SEP on later life immune response to CMV, adjusted for age and gender, N=1562



Supplemental Figure 2. Diagram of SEM for standardized direct and indirect effects of early life SEP on later life immune response to *T. gondii*, adjusted for age and gender, N=1562




Supplemental Figure 3. Diagram of SEM for standardized direct and indirect effects of early life SEP on later life immune response to *H. pylori*, adjusted for age and gender, N=1562




\* $p < 0.05$ , \*\* $p < 0.01$

Supplemental Figure 4. Log Odds (SE) for SEM Model of Low v. High Cumulative Immune Response adjusted for age and gender, N= 1263



Total Effect of ChildSEP on Low v. High Cumulative Immune Response: 0.16 (0.08)  
 Total Indirect Effect of ChildSEP on Low v. High Cumulative Immune Response: 0.11 (0.05)\*  
 Indirect Effect via Education Pathway: 0.09 (0.05)\*  
 Indirect Effect via Education-Occupation Pathway: 0.02 (0.04)

Supplemental Figure 5. Log Odds (SE) for SEM Model of Middle v. High Cumulative Immune Response adjusted for age and gender, N= 1263



Total Effect of ChildSEP on Middle v. High Cumulative Immune Response : -0.20 (0.12)  
 Total Indirect Effect of ChildSEP on Middle v. High Cumulative Immune Response: -0.12 (0.06)  
 Indirect Effect via Education Pathway: -0.09 (0.06)  
 Indirect Effect via Education-Occupation Pathway: -0.03 (0.02)

† Modeling Middle v. High Cumulative Immune Response Class  
 \* Significant at p=0.05; \*\* Significant at p=0.01;\*\*\* Significant at p=0.001